



READ SAFETY DIRECTIONS BEFORE OPENING OR USING

# ROUT<sup>®</sup>

*Ornamental  
Herbicide*

**Active Constituents:** 20g/kg Oxyfluorfen  
10g/kg Oryzalin



20kg  
Net Weight





20kg  
Net Weight



20kg  
Net Weight

# ROUT<sup>®</sup>

## Ornamental Herbicide

### DIRECTIONS FOR USE (In all states)

#### Restraints

- Do NOT mix ROUT<sup>®</sup> into potting media as poor weed control will be obtained and plant damage is likely.
- Do NOT apply ROUT<sup>®</sup> in wind conditions likely to affect good distribution.
- Do NOT broadcast ROUT<sup>®</sup> over wet foliage.

Situation	Weeds Controlled	Application Rate	Critical Comments
Container ornament al plants (See table below)	Weeds controlled by ROUT <sup>®</sup> include the following: Barley grass Barnyard grass Burr grass Crowsfoot grass Liverseed grass Love grass Pigeon grass Rhodes grass Rye grass Stink grass Summer grass Winter grass Amsinckia Bladder ketmia Blackberry nightshade Caltrop, Capeweed, Chickweed, Creeping oxalis, Deadnettle, Fat hen, Fleabane, Flickweed, Giant pigweed, Mallow, Mimosa, Pigweed, Prickly Lettuce, Red caustic creeper, Redshank (Amaranth), Shepards purse, Soursob, Sowthistle, Staggerweed, Starrburr, Subterranean clover, Thornapple, White eye, Wild mustard, Wild radish, Wireweed, Willowherb	100 kg/ha or 1.0kg/100 square metres or 10g/square metre	Apply evenly over the area or pots to be treated preferably using a properly calibrated mechanical granule spreader. Apply to the surface of weed free soil preferably after cultivation or just after potting up. ROUT <sup>®</sup> will not control established weeds. Apply to dry foliage only and then sprinkler irrigate or hose (10-15mm) to wash granules from foliage and incorporate the herbicide into the soil.  Do not apply ROUT <sup>®</sup> to wet foliage - should this happen irrigate or hose immediately to remove granules from the leaf surface.  Follow the precautions listed below to avoid unnecessary plant damage and to obtain optimum weed control.  One application will last for 2-3 months depending upon rainfall, irrigation, soil type (potting mix), temperature and weed seed burden. Retreat when the first signs of weed growth are seen but no sooner than 2 months and no later than 3 months after the first application.  Do not apply ROUT <sup>®</sup> 2 weeks prior to or 2 weeks after leaf bud break or during periods of flush growth. Newly formed leaves are sensitive to this product. Spotting or leaf drop may result from such applications.  Do not apply directly onto bare roots of woody ornamentals as injury may occur.

**NOT TO BE USED FOR ANY PURPOSE OR IN ANY MANNER CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.**

**IN TASMANIA THIS PRODUCT MUST NOT BE APPLIED BY AIRCRAFT WITHOUT SPECIFIC APPROVAL OF THE REGISTRAR.**

#### GENERAL INSTRUCTIONS

**Ornamentals on which ROUT<sup>®</sup> may be used**

**Australian testing has indicated that the following species are tolerant of ROUT<sup>®</sup> when used as recommended:**

Abelia x grandiflora 'FrancisMason' (Glossy abelia), Acer palmatum (Japanese maple), Acacia baileyana (Cootamundra wattle), Acacia acinacea, Alnus acuminata, Arbutus unedo (Strawberry tree), Azalea indica, Betula pendula (Birch), Bougainvillea spp., Buxus microphylla v. japonica (Japanese box), Callistemon viminalis 'Captain Cook' (Bottlebrush), Camellia japonica, Cassia bicapsularis, Casuarina littoralis, Casuarina torulosa, Choisya ternata (Mexican orange), Coleonema album, Cupressus sempervirens v. (Swanes golden pencil pine), Eriostemon myoporoides, Eucalyptus nicholi (Fine leaf peppermint), Eucalyptus sideroxylon (Pink iron bark), Fraxinus excelsior (Golden ash), Fraxinus pennsylvanica (Red ash), Gardenia jasminoides, Gardenia magnifica, Gelsemium sempervirens (Carolina jasmine), Gordonia axillaris, Grevillea 'Clearview David', Grevillea robusta (Silky oak), Grevillea arenaria, Grevillea 'Poorinda', Hakea saligna, Hakea laurina, Hebes buxifolia, Hibiscus sp., Juniperus conferta, Leptospermum petersonii, Leptospermum juniperinum 'Horizontalis', Melaleuca armillaris, Melaleuca decussata, Melia azedarach, Murraya exotica (orange jasmine), Nandina domestica 'Nana' (Dwarf nandina), Nerium oleander (Oleander), Photinia robusta, Punica granatum (Pomegranate), Prunus sp. (Weeping cherry), Rosa sp. (Rose), Rosmarinus folia ('Blue lagoon' rosemary), Syringa (Lilac), Ulmus procera (English elm), Ulmus campestris 'Variegata' (Silver elm sp.), Westringia fruticosa.

**Overseas testing under similar conditions to Australia has indicated that the following species are tolerant of ROUT<sup>®</sup> when used as recommended:**

Cassia corymbosa, Cotoneaster sp., Diosma ericoides, Euonymus alata, Ficus benjamina, Iberis sempervirens, Ilex vomitoria, Jasminium nitidum, Juniperus chinensis, Juniperus sabina, Juniperus squamata, Lagerstroemia indica, Lantana montevidensis, Ligustrum japonicum, Ligustrum lucidum, Ligustrum sinense, Picea abies (Norway Spruce), Picea glauca (Dwarf Alberta Spruce), Pinus elliotii (Slash pine), Pinus radiata, Pinus strobus, Pinus thunbergiana, Pittosporum crassifolium, Pittosporum tenuifolium, Pittosporum tobira, Podocarpus gracilor, Raphiolepis indica, Rhododendron spp.(Azalea), Schefflera arboricola, Thuja occidentalis, Viburnum plicatum, Weigela spp.

**Ornamentals on which ROUT<sup>®</sup> Ornamental Herbicide should not be used:**

**Overseas and local testing has indicated that the following species can be sensitive to ROUT<sup>®</sup>:**

Dietes bicolor, Euonymus fortunei, Euonymus japonicus, Europs pectinatus, Ficus pumila (Creeping fig), Hardenbergia spp., Ilex comuta, Potentilla fruticosa, Salvia farinacea, Trachelspermum asiaticum.  
**ROUT<sup>®</sup> IS NOT RECOMMENDED FOR USE ON VEGETABLES OR EDIBLE FRUIT CROPS, TURF, BEDDING PLANTS OR GREENHOUSE FLOWERING CROPS.**

#### APPLICATION OF ROUT<sup>®</sup>

ROUT may be applied individually to each pot or pots can be grouped together and ROUT distributed using a granule applicator or GLOVED hand.

Ensure ROUT does not stick to plant foliage or remain lodged in leaf axils or other plant structures.

To assist in application/calibration 100 kg/ha of ROUT is equivalent to 30/32 granules per 10 square centimetres (2 cm x 5 cm). Even application is important for good weed control. DO NOT disturb the soil surface once the product has been applied as the herbicide barrier will be broken and weed growth can occur.

#### PRECAUTIONS

DO NOT apply to pots within 2 weeks of retail sale. DO NOT apply in enclosed glasshouse type structures or within 2 weeks prior to enclosure in such structures.

#### Protection of crops, native and other non-target plants

DO NOT apply to whorl leaved plants as their leaves channel the herbicide granules to the leaf base making removal of the granules difficult. DO NOT apply under weather conditions, or from equipment, that may cause drift onto susceptible plants/crops, cropping lands or pastures.

#### Protection of livestock

DO NOT graze any treated plant or cut for stockfood or feed to poultry.  
DO NOT contaminate water used by livestock.

#### Protection of wildlife, fish, crustacea and environment

Dangerous to wildlife, fish and other aquatic organisms. DO NOT contaminate streams, rivers or waterways with the product or used container, or when cleaning equipment. DO NOT use product where fauna are foraging and are likely to feed on granules.

#### Spills

Sweep up spilled material and return to original container. Wash contaminated equipment taking care that washings do not enter sewers, drains and other waterways. Dispose of waste according to the Australian Standard AS 2507 - Storage and Handling of Pesticides. DO NOT allow spilled product to enter sewers, drains, creeks or any other waterways.

#### Storage and disposal

Keep out of reach of children. Store in a locked room or place away from children, animals, food, feedstuffs, seed and fertiliser. Store in the closed original container, in a dry, cool well-ventilated area out of direct sunlight. Shake containers empty onto treatment site. Puncture or shred and bury empty bags in a local authority landfill. If no landfill is available, bury the containers below 500mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty bags and product should not be burnt.

#### SAFETY DIRECTIONS

Avoid contact with the eyes and skin. Do not inhale dust, use a dust mask. If product on skin immediately wash the area with soap and water. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water.

#### FIRST AID

If poisoning occurs contact a doctor or Poisons Information Centre (Phone 13 11 26).

#### MATERIAL SAFETY DATA SHEET

Further information on this product is contained in a Material Safety Data Sheet (MSDS) available from product suppliers or the manufacturer.

#### EMERGENCY INFORMATION PANEL

<b>DANGEROUS GOOD CLASS</b>	The product is not classified as a Dangerous Good by the Australian ADG Code
<b>IN EMERGENCY DIAL 000 POLICE OR FIRE BRIGADE</b>	<b>SPECIALIST ADVICE IN EMERGENCY ONLY ALL HOURS 1800 033 111</b>

ROUT<sup>®</sup> is a registered Trademark of The Scotts Company and it's affiliates.

APVMA No. 31548/20/1103

# NET CONTENTS: 20kg



**Scotts Australia Pty. Ltd**

11 Columbia Way, Baulkham Hills, New South Wales, 2153  
Phone 02 8853 7300 • Fax 02 8853 7310  
www.scottsaustralia.com

Rout<sup>®</sup> is the registered trademark of The Scotts Company and it's affiliates.