

CAUTION

KEEP OUT OF REACH OF CHILDREN
 READ SAFETY DIRECTIONS BEFORE OPENING OR USING

DICOT III 21:1:16**FERTILISER WITH BROADLEAF HERBICIDE**

Herbicide Active Constituents:

10.3 g/kg	2,4-D ACID
10.3 g/kg	MECOPROP ACID
0.7 g/kg	DICAMBA ACID

Guaranteed Fertiliser Analysis:

TOTAL NITROGEN (N)	21.00%
As ammonium	0.5%
As urea and methylene ureas	20.5%
As water soluble organic nitrogen	8.8%
As water insoluble, nitrogen	1.4%
PHOSPHORUS (P)	1.3%
POTASSIUM (K)	16.6%
SULPHUR (S) as sulphate	7.5%

GROUP I HERBICIDE**COMBINATION HERBICIDE AND NPK FERTILISER**

- **FOR POST EMERGENT CONTROL OF BROADLEAF WEEDS & FEEDING OF TURF WITH SLOW RELEASE NITROGEN**
 - **NOT FOR USE ON BENT GRASS GREENS, BUFFALO GRASS, CARPET GRASS OR DICHONDRA.**

DIRECTIONS FOR USE

RESTRAINTS: DO NOT apply if rain or irrigation is expected within 24 hours. DO NOT apply when daytime temperatures are consistently above 32°C. DO NOT apply to newly established turf until 4 weeks after the first mowing. DO NOT plant grass seed until 1 month after application. DO NOT over apply, especially under the branch spread of trees and shrubs. DO NOT use on bent grass, Buffalo grass, carpet grass or Dichondra. DO NOT apply if weeds are stressed due to dry, frosty or excessively moist conditions.

SITUATION	TARGET	RATE	CRITICAL COMMENTS
Turf	Post emergent control of broadleaf weeds including bindii, capeweed, Catsear (flatweed), chickweed, mouse-eared chickweed, clovers, dandelion, fat hen, creeping oxalis, mallow, pearlwort, hydrocotyl, plantain (lambs tongue), stagger weed, fumitory, cobblers peg and wireweed.	160kg/ha (1.6kg/ 100m ²)	The product is ready to apply. No mixing or dilution is required. Apply to moist foliage using a granular spreader able to give even coverage. If foliage is not moist from dew or recent rain sprinkler irrigate the area to be treated. Rain, sprinkling, mowing or heavy traffic within 24 hours can knock granules off weeds and may reduce effectiveness, although the herbicides in the product are both foliar and root absorbed. Retreating may be necessary, but wait at least three weeks to observe the full results of the first application. For best results do not mow turf grass for 48 hours after application of this product.
	Turf nutrition - Provides slow release nitrogen as methylene urea for 6- 10 weeks feeding		

NOT TO BE USED FOR ANY PURPOSE OR IN ANY MANNER CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION

GENERAL INSTRUCTIONS:

DICOT III 21:1:16 is a convenient ready to use formulation for the control of many broadleaf weeds in turf and also for fertilisation of turf. The product should only be used when both functions are required.

For application use a Scotts R8-A professional spreader or an equivalent granular spreader which can be calibrated for even output and capable of giving even application. Apply under calm conditions, thereby reducing drift to non-target areas and optimising particle distribution for good weed control. Best results will be obtained when weeds are actively growing.

Warning: Do not use on or in situations where damage to susceptible crops such as cotton, vines tomatoes, vegetables, fruit trees, lucerne, legume crops and pastures, oilseed crops or other susceptible crops and trees (eg. Kurrajongs, belahs and eucalypts) may occur from direct application, drift including storm run-off or vapour.

Resistant Weeds Warning: DICOT III 21:1:16 herbicide is a member of the benzoic acid and phenoxy Group of herbicides. The product has the disruptor of plant cell growth mode of action. For weed resistance management, the product is a Group I herbicide. Some naturally occurring weed biotypes resistant to the herbicide may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds will not be controlled by this product or other disruptor of plant cell growth herbicides. Since the occurrence of resistant weeds is difficult to detect prior to use, Scotts Australia Pty. Ltd. accepts no liability for any losses that may result from failure of this product to control resistant weeds.

SETTINGS TO APPLY RECOMMENDED RATE					
	SCOTTS DROP	SCOTTS ROTARIES (R-8A SPREADER)	ACCUPRO 2000	VICON MODEL 02	VICON MODEL 03
Ground speed	4.5km/hr	4.5km/hr	4.5km/hr	7km/hr	7km/hr
Width of coverage	Overlap	3.65m	3.65m	9.15m	7.3m
Cone Setting		8	8		
Half rate/double application	23	K½	K½	15	18
Normal rate/single application	22	J	J	23	25

PROTECTION OF WILDLIFE, FISH, CRUSTACEA AND ENVIRONMENT: Do not contaminate dams, streams or other waterways with the product or used containers.

SPILLS: Sweep up spilled material and return to original container. Wash contaminated equipment taking care that washings do not enter sewers, drains and other waterways.

STORAGE AND DISPOSAL: Store in the closed original container, in a dry, cool, well ventilated area out of direct sunlight. Do not store with weed, feed, foodstuffs, insecticides, fungicides or other fertilisers or herbicides. Puncture or shred empty container and bury in a local authority landfill. If no landfill is available bury the containers below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation or tree roots. Empty containers or product should not be burnt. Do not expose of excess or damaged material on site.

SAFETY DIRECTIONS: Will irritate the eyes. Avoid contact with eyes. When opening the container and using the product, wear elbow length PVC gloves goggles and disposable dust mask. If product in eyes, wash it out immediately with water. Wash hands after use. After each day's use, wash gloves and goggles.

FIRST AID: If poisoning occurs, contact a doctor or Poisons Information Centre [Ph. 13 11 261. If skin contact occurs, remove contaminated clothing and wash skin thoroughly.

If in eyes, hold eyes open, flood with water for at 1-15 minutes and see a doctor.

IMPORTANT NOTICE LIMITATION OF LIABILITY. This product has been researched to provide necessary data to support its uses listed on the label. The user should always follow the label directions and exercise his/her judgement and caution when using the product under his/her growing conditions. **NO WARRANTY OR REPRESENTATION IS MADE, EXPRESSED OR IMPLIED, CONCERNING THE RESULTS OBTAINED FROM THE USE OF THIS PRODUCT IF NOT IN ACCORDANCE WITH DIRECTIONS OR ESTABLISHED SAFE PRACTICE.** The exclusive remedy of the user or buyer, and the limit of liability of The Scotts Company or its affiliates, for any and all losses, injuries or damages resulting from the use or handling of this product shall be the purchase price paid by the user or buyer for the quantity of this product involved. The buyer and users are deemed to have accepted the terms of this Notice, which may be varied only by agreement in writing, signed by a duly authorised representative of The Scotts Company.

NET CONTENTS: 20 kg

EMERGENCY INFORMATION PANEL

IN EMERGENCY DIAL 000, POLICE OR FIRE BRIGADE	SPECIALIST ADVICE IN AN EMERGENCY ONLY 1800 033 111
--	--

MSDS: Further information on this product is contained in the Material Safety Data Sheet, copies of which are available from product suppliers or the manufacturer.

**Scotts Australia Pty. Ltd. ABN. 31 003 123 162. 11 Columbia Way, Baulkham Hills, NSW, 2153.
Phone (02) 8853 7300 Fax: (02) 8853 7310. APVMA APPROVAL NO: 56198/1202**

8035

As separate sticker on side of container

Batch Number		Date of Manufacture	
--------------	--	---------------------	--